



ACC.14

TCT@ACC-12 | innovation in intervention

A1187

JACC April 1, 2014

Volume 63, Issue 12



Non Invasive Imaging

COMBINED ATRIOVENTRICULAR LONGITUDINAL STRAIN RATE DURING ISOVOLUMIC CONTRACTION PREDICTS PULMONARY CAPILLARY WEDGE PRESSURE IN PATIENTS WITH SYSTOLIC DYSFUNCTION

Poster Contributions

Hall C

Sunday, March 30, 2014, 3:45 p.m.-4:30 p.m.

Session Title: Non Invasive Imaging: Myocardial Strain, Cardiac Mechanics and Diastolic Function

Abstract Category: 15. Non Invasive Imaging: Echo

Presentation Number: 1210-42

Authors: *Alaa M. Omar, Mohamed A. Abdel-Rahman, Mostafa Helmy, Hazem Khorshid, Hala Raslan, Osama Rifaie, National Research Centre, Medical Division, Department of Internal Medicine, Cairo, Egypt, Ain Shams University Hospital, Department of Cardiology, Cairo, Egypt*

Background: We recently reported that tissue Doppler (TD) mitral annular isovolumic contraction (IC) velocity (IVV) predicts pulmonary capillary wedge pressure (PCWP) in systolic dysfunction. Aim: Study the ability of TD-derived longitudinal left atrial (LA) and left ventricular (LV) strain rate (SR) during IC in predicting PCWP.

Methods: Thirty-five patients with heart failure symptoms were studied (age: 56 ± 8 years, 12 (35%) females, and EF: $51 \pm 14\%$). PCWP was invasively measured immediately after echocardiographic measurement of IVV and peak longitudinal systolic LV and LA SR during IC and ejection (EJ). LV-SR were calculated from basal and mid segments of LV walls in apical views, while mean LA-SR during IC and EJ, and IVV were calculated from 4 LA walls, and their corresponding mitral annular positions. Atrioventricular SR (AV-SR) during IC and EJ was calculated as the sum of LV and LA values. Patients were then classified into 17 (49%) with EF $\geq 55\%$, and 18 (51%) with EF $< 55\%$ (62.7 ± 4.4 vs. 40.2 ± 10 , $p < 0.001$).

Results: Age, Sex, risk factors, and medications were similar between both groups. In EF $< 55\%$, LV-SRI during EJ, LA-SRI during IC and EJ, and IVV were lower (1 ± 0.3 vs. 1.2 ± 0.2 , 1.3 ± 0.6 vs. 1.9 ± 1 , 1.5 ± 0.5 vs. 2.6 ± 1.3 s⁻¹, 4.63 ± 1.2 vs. 7.01 ± 1.9 cm/s, $p = 0.03$, 0.03 , 0.001 , < 0.001). AV-SRI during IC and EJ were also lower in patients with EF $< 55\%$ (2 ± 0.8 vs. 2.7 ± 1.06 s⁻¹, 2.5 ± 0.6 vs. 3.9 ± 1.1 s⁻¹, $p = 0.023$, < 0.001). LV-SRI during IC and EJ, LA-SRI during IC, AV-SRI during IC, and IVV correlated with PCWP in EF $< 55\%$ ($r = -0.59$, -0.55 , -0.68 , -0.72 , -0.63 , $p = 0.01$, 0.02 , 0.002 , < 0.001 , 0.005). Other correlates with PCWP in EF $< 55\%$ were E/e' and left atrial volume ($r = 0.47$, 0.7 , $p = 0.04$, 0.001 ; respectively). Multivariate regression in EF $< 55\%$ revealed that AV-SRI-IC was the only independent predictor of PCWP. Finally, IVV correlated with LV and LA SRI-IC and this correlation became strongest with AV-SRI during IC ($r = 0.65$, 0.71 , 0.77 , $p = 0.01$, 0.002 , 0.001 ; respectively).

Conclusion: in our study we showed that LA and LV SRI-IC act mutually to correlate with PCWP in patients with systolic dysfunction. The correlation between mitral annular IVV and PCWP in those patients might be a product of this combination.